

Effects of amyloid- β peptide on glutamine transporter mRNA expression and cell viability in cultured rat cortical cells

Doungjai Buntup^a, Anek Chayasadam^b, Rudee Surarit^c, Nuanchan Jutapakdeegul^a, Wipawan Thangnipon^{a,*}

^a Neuro-Behavioural Biology Centre, Institute of Science and Technology for Research and Development, Mahidol University, Nakorn Pathom 73170, Thailand

^b Department of Periodontology, Faculty of Dentistry, Khon Kaen University, Khon Kaen 40002, Thailand

^c Department of Physiology and Biochemistry, Faculty of Dentistry, Mahidol University, Bangkok 10400, Thailand

*Corresponding author, e-mail: grwtn@mahidol.ac.th

Received 11 Nov 2008

Accepted 27 May 2009

ABSTRACT: Alzheimer's disease is a major neurodegenerative disorder in which there is an overproduction and accumulation of amyloid- β ($A\beta$) peptides. During the initial stages of the disease, glutamate receptors are dysregulated by $A\beta$ accumulation resulting in the disruption of glutamatergic synaptic transmission. We used rat cortical cell cultures to examine the effects of $A\beta(25-35)$ -induced neurotoxicity on glutamine transporters involved in the glutamate cycle. In primary mixed cell cultures prepared from cerebral cortex, incubation with 10 μ M $A\beta(25-35)$ for 12 h, but not for 24 h, markedly suppressed system A transporter 1 (SAT1) mRNA expression. On the other hand, $A\beta(25-35)$ had no effect on SAT1 mRNA level in neuronal cell cultures. Treatment of both types of cell cultures with $A\beta(25-35)$ resulted in a significant decrease in cell survival in a concentration and time-dependent manner, as determined by MTT assay. These results indicated that $A\beta$ may impair neuronal function and transmitter synthesis and perhaps reduce excitotoxicity through a reduction in neuronal glutamine uptake.

KEYWORDS: Alzheimer's disease, amyloid- β peptide, mRNA glutamine transporter

INTRODUCTION

Alzheimer's disease is a neurodegenerative disorder characterized by the presence of neurofibrillary tangles, neuritic plaques, synaptic depletion, and neuronal cell loss in specific cortical and subcortical areas of the brain¹. Senile plaques contain amyloid- β ($A\beta$) peptide, produced by cleavage of the amyloid precursor protein (APP) by β - and γ -secretases². $A\beta$ is a peptide of 39–43 amino acids that forms insoluble aggregates surrounded by degenerating neuritis and activated glial cells³. Numerous experiments with synthetic $A\beta(1-42)$ suggest that this peptide is toxic to cultured neurons by inducing protein oxidation, lipid peroxidation, and oxidative stress in the cellular environment in ways that are inhibited by antioxidants of free-radicals⁴. In culture, $A\beta$ can directly induce neuronal cell death⁵ and can render neurons vulnerable to excitotoxicity⁶ and oxidative insults⁷.

Glutamate is the major excitatory neurotransmitter of the mammalian central nervous system that contributes not only to fast synaptic neurotransmission

but also to complex physiological processes, such as memory, learning, plasticity, and neuronal cell death⁸. The neurons use glutamine as the main substrate for synthesis of glutamate and γ -aminobutyric acid^{9,10}. The release of glutamine from astrocytes and the uptake of glutamine by neurons are integral steps in the glutamate-glutamine cycle, a major pathway for the replenishment of neuronal glutamate¹¹.

Recently, cDNA encoding three distinct system A amino acid transporters (SAT1 to SAT3) have been cloned and functionally identified¹². SAT1 isoform is preferentially expressed on the plasma membrane of glutamatergic neurons and encodes for a highly efficient glutamine transporter^{13,14}. Dolinska et al¹⁵ reported that SAT1 mRNA is transcribed in neurons but not in cerebral cortical astrocytes or cerebellar astrocytes. A malfunction in the glutamate transport system can lead to accumulation of excessive glutamate in the synapse, which is harmful to neurons and could result in neurodegeneration^{16,17}. However, the causal relationship between $A\beta$ and glutamine transporters in Alzheimer's disease is not yet well

defined.

A small 11-amino acid fragment of the full-length peptide, A β (25–35), is a convenient alternative in Alzheimer's disease investigations as the peptide mimics several toxicological and oxidative stress properties of the native full-length peptide^{4,18,19}. Recently, our group demonstrated that treatment of mixed cell cultures with 10 μ M A β (25–35) for 12 h and 24 h reduces SAT1 protein level as demonstrated by immunocytochemical and quantitative immunoreactivity analysis²⁰. We now report the effects of A β (25–35) on SAT1 expression in rat cortical primary cell cultures.

MATERIALS AND METHODS

Enriched primary neuronal cell cultures were prepared from cerebral cortices of 10 embryonic day-18 (E18) Wistar rat fetuses as described previously^{21,22}. Cells plated at 5×10^5 cells/ml were cultured in poly-L-lysine coated multiwell plates and maintained in serum-free Dulbecco's modified Eagles medium (DMEM) supplemented with B27 under a humidified atmosphere of 5% CO₂/95% air at 37 °C for 5 days before incubating with peptide.

Primary mixed cell cultures were prepared from cerebral cortices of 10 1-day old Wistar rat neonates as previously described²³. Cells plated at 5×10^5 cells/ml were cultured in poly-L-lysine coated multiwell plates. The dissociated cortical cells were suspended in Neurobasal-A medium containing 2% (v/v) B27 and 1% (v/v) GlutaMAX I and maintained as described for the E18 cells.

The B27-containing medium was removed from neuronal and mixed cell cultures at day 6. Cells were then washed twice with DMEM or Neurobasal-A medium and then incubated for 12 h and 24 h in the absence or presence of 3 and 10 μ M A β (25–35) (Sigma) to produce a sub-lethal neurotoxicity.

Total RNA was isolated using RNeasy Mini Kit (Qiagen) and the yield was determined by spectrophotometry at 260 nm. The Reverse transcriptase polymerase chain reaction (RT-PCR) was carried out using the Qiagen one step RT-PCR kit. PCR amplifications of SAT1 employed primers 5'-ACAGGC-GACATTCTCATCCT-3' (forward) and 5'-GTTTCA-GTGGCCTTACCAT-3' (reverse), giving rise to a 426 base pair amplicon²⁴, and β -actin (housekeeping gene) primers 5'-CCCAGAGCAAGAGAGGCATC-3' (forward) and 5'-CTCAGGAGGAGCAATGATCT-3' (reverse), giving rise to a 830 base pair amplicon²⁵. Primers were obtained from the Bioservice Unit of the National Science and Technology Development Agency, Thailand. Thermocycling conditions, opti-

mized in order to allow amplicon synthesis within the linear-log phase of amplification, were as follows: reverse-transcription at 50 °C for 30 min; inactivation of reverse transcriptase and activation of Hot start *Taq* DNA polymerase at 95 °C for 15 min; 30 cycles of 30 sec at 94 °C, 1 min at 55 °C (for STAT1) or 1 min at 72 °C (for β -actin); and a final extension step at 72 °C for 10 min. Amplicons were separated by electrophoresis in 2% agarose gel and stained with ethidium bromide. Densitometric analysis was performed using QUANTITY ONE 4.5.1 (Bio-Rad).

This method is based on the reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) tetrazolium salt to a crystalline blue formazan product by cellular oxidoreductase²⁶. Therefore, the amount of formazan produced is proportional to the number of viable cells. The culture medium was removed from the treated cells and replaced with a solution of MTT (0.5 mg/ml) in phosphate-buffered saline, pH 7.4. After incubation for 4 h at 37 °C, the solution was removed and the resulting blue formazan solubilized in 0.1 ml of 0.04 M HCl in isopropanol. Absorbance at 570 nm was measured at 630 nm using a microplate reader (Bio-Tex EL 311).

Data were analysed by one-way ANOVA using LSD for post-hoc comparisons. Differences were considered statistically significant if $P < 0.05$. Each experiment was repeated 3–4 times and results from a representative experiment are shown.

RESULTS

To determine the effect of A β (25–35) on expression of SAT1 in both neuronal and mixed cell cultures, cells were cultured for 5 days and then incubated with A β (25–35) for 12 h and 24 h. SAT1 mRNA levels were determined relative to those of β -actin²⁷.

The RT-PCR of the mixed cell culture showed that 10 μ M A β (25–35) significantly suppressed SAT1 mRNA expression after incubation for 12 h ($P < 0.001$) (Fig. 1). However, SAT1 expression at 24 h showed no variation after exposure to 3 and 10 μ M of A β (25–35). In neuronal cell culture, expression of SAT1 mRNA did not change after exposure to 3 and 10 μ M of A β (25–35) for 12 and 24 h when compared with the control (Fig. 2).

Neuronal cell cultures showed a significant loss of cell viability, as determined by MTT method, when treated for 24 h with A β (25–35) in a dose-dependent manner ($P < 0.05$) (Fig. 3). A significant cell loss was also observed after treatment with 10 μ M A β (25–35) for 12 h ($P < 0.05$). Viability of mixed cell cultures was reduced when exposed to 3 and 10 μ M




Fig. 1 Expression of SAT1 mRNA in mixed cell culture exposed to Aβ(25–35). Mixed cell cultures were incubated with Aβ(25–35) for 12 and 24 h and subjected to RT-PCR for amplification of SAT1 and β-actin mRNA. (a) Amplicons separated by gel-electrophoresis and stained with ethidium-bromide (b) quantitated bands. Lane 1: control; lane 2: exposed to 3 μM Aβ(25–35); lane 3: exposed to 10 μM Aβ(25–35). ** $P < 0.001$ compared with control. Columns show mean \pm SEM from three independent experiments.




Fig. 2 Expression of SAT1 mRNA in the neuronal cell culture exposed to Aβ(25–35). Neuronal cell cultures were incubated with Aβ(25–35) for 12 and 24 h and subjected to RT-PCR for amplification of SAT1 and β-actin mRNA. (a) Amplicons separated by gel-electrophoresis and stained with ethidium-bromide (b) quantitated bands. Lane 1: control; lane 2: exposed to 3 μM Aβ(25–35); lane 3: exposed to 10 μM Aβ(25–35). Columns show mean \pm SEM from three independent experiments.

Aβ(25–35) for 12 and 24 h ($P < 0.05$ and $P < 0.001$ respectively).

DISCUSSION

Results from this study indicated a significant decrease of SAT1 mRNA level after 12 h exposure to 10 μM Aβ(25–35) in mixed but not in neuronal cell cultures. Baron et al²⁸ suggested that Aβ exerts its toxic effect via activation of transcription factors. Aβ(25–35) induces expression of the growth arrest and DNA damage-inducible gene (gadd 45) implicated in the DNA excision-repair process. Accumulation and oligomerization of Aβ is also thought to play a central role in Alzheimer's disease pathogenesis. Aβ can produce oxidative damage by stimulating neuroinflammation, as well as generating reactive oxygen species as a result of binding to mitochondrial proteins²⁹. Studies of neuron co-cultures showed that astrocyte formation of nitric oxide and other reactive oxygen species does occur in a number of neurodegenerative disorders^{30,31}. Thus a mixed cell culture will be more sensitive to toxic effects than the neuronal cell culture when exposed to Aβ. Our data imply that a decrease of glutamine uptake may play

an important role in Alzheimer's disease progression.

However, Aβ(25–35) had no effect on SAT1 mRNA expression at 24 h in both mixed and neuronal cell cultures. Tian et al³² showed that SAT2 mRNA level does not decrease after exposure to a low dose of Aβ(25–35). This effect may act as a cellular defence against neurodegeneration in pathogenesis of Alzheimer's disease.

Viability of mixed and neuronal cell cultures, measured by MTT method, was reduced when exposed to Aβ(25–35) in both a time- and concentration-dependent manner. In neuron-enriched cultures, Aβ is known to increase vulnerability to excitotoxicity³³. Wei et al³⁴ showed that Aβ(1–42), in a time (1–48 h) and concentration (0.01–20 μM)-dependent manner, induces toxicity in cultured neurons. Similarly, Domenici et al³⁵ demonstrated that while pure neuronal culture shows a significant cell loss only at the highest concentration of Aβ(25–35), mixed cell culture manifests a toxic effect in a dose-dependent manner which was significant down to the lowest concentration. As observed by Pfrieger and Barres³⁶, neurons co-cultured with astrocytes develop




Fig. 3 Effect of A β (25–35) on neuronal cell viability. (a) Mixed and (b) neuronal cell cultures were treated with 3 and 10 μ M A β (25–35) for 12 and 24 h and cell viability was determined using MTT assay. * $P < 0.05$ and ** $P < 0.001$ compared to control. Data show mean \pm SEM of four independent experiments.

approximately 7-fold more synapses and have a 7-fold increase in synaptic efficacy compared with neurons raised in the absence of astrocytes. Astrocytes could contribute to neurodegeneration, as A β -activated astrocytes overexpress factors such as interleukin 1 β , nitric oxide, and S100 β ^{37–39}, which are all potentially neurotoxic³⁵. In summary, our findings show that the reduction of cell viability is more prominent in mixed cell cultures ($P < 0.001$) than neuronal cell cultures ($P < 0.05$) when exposed to A β . Additionally, in mixed cell cultures, A β markedly suppressed SAT1 mRNA expression ($P < 0.001$) but not in cultured neuronal cells. This observation supports our previous report²⁰ that the presence of astroglia enhances A β induced neurotoxicity in cultured cortical cells. A possible explanation is that neurons can be damaged indirectly by the effect of A β -astrocyte interaction. In addition, short-time exposure to 0.1–50 μ M A β (25–35) is not toxic to neurons, while longer exposures to the peptide are increasingly toxic⁴⁰. Recently, Geci et al⁴¹ showed that incubation of neuronal cell cultures with lower concentrations of A β (25–35) (5 and 10 μ M) trigger apoptosis within 24 h of treatment. **Our study suggests that A β may impair neuronal function and transmitter synthesis, and perhaps re-**

duce excitotoxicity through a reduction in neuronal glutamine uptake.

Acknowledgements: This work was supported by the Thailand Research Fund through the RGJ PhD Programme (PHD/0230/2545). The animal procedures were approved by the committee on ethics in animal research of the Institute of Science and Technology for Research and Development, Mahidol University, MUSTA 2006/003.

REFERENCES

- D'Aniello A, Fisher G, Migliaccio N, Cammisà G, D'Aniello E, Spinelli P (2005) Amino acids and transaminases activity in ventricular CSF and in brain of normal and Alzheimer patients. *Neurosci Lett* **388**, 49–53.
- Govaerts L, Schoenen J, Bouhy D (2007) Pathogénie de la maladie d'alzheimer: les mécanismes moléculaires et cellulaires. *Rev Med Liege* **62**, 209–16.
- Abramov AY, Canevari L, Duchon MR (2004) Calcium signals induced by amyloid- β peptide and their consequences in neurons and astrocytes in culture. *Biochim Biophys Acta* **1742**, 81–7.
- Varadarajan S, Kanski J, Aksenova M, Lauderback C, Butterfield DA (2001) Different mechanisms of oxidative stress and neurotoxicity for Alzheimer's A β (1–42) and A β (25–35). *J Am Chem Soc* **123**, 5625–31.
- Ueda K, Fukui Y, Kageyama H (1994) Amyloid β protein-induced neuronal cell death: neurotoxic properties of aggregated amyloid β protein. *Brain Res* **639**, 240–4.
- Koh JY, Yang LL, Cotman CW (1990) β -amyloid protein increases the vulnerability of cultured cortical neurons to excitotoxic damage. *Brain Res* **533**, 315–20.
- Goodman Y, Mattson MP (1994) Secreted forms of β -amyloid precursor protein protect hippocampal neurons against amyloid β -peptide-induced oxidative injury. *Exp Neurol* **128**, 1–12.
- Shigeri Y, Seal RP, Shimamoto K (2004) Molecular pharmacology of glutamate transporters, EAATs and VGLUTs. *Brain Res Rev* **45**, 250–65.
- Erecińska M, Silver IA (1990) Metabolism and role of glutamate in mammalian brain. *Progr Neurobiol* **35**, 245–96.
- Struzyńska L, Sulkowski G (2004) Relationships between glutamine, glutamate, and GABA in nerve endings under Pb-toxicity conditions. *J Inorg Biochem* **98**, 951–8.
- Broer S, Brookes N (2001) Transfer of glutamine between astrocytes and neurons. *J Neurochem* **77**, 705–19.
- Reyna SV, Ensenat D, Johnson FK, Wang H, Schafer AI, Durante W (2004) Cyclic strain stimulates L-proline transport in vascular smooth muscle cells. *Am J Hypertens* **17**, 712–7.
- Varoqui H, Zhu H, Yao D, Ming H, Erickson JD (2000)

- Cloning and functional identification of a neuronal glutamine transporter. *J Biol Chem* **275**, 4049–54.
14. Armano S, Coco S, Bacci A, Pravettoni E, Schenk U, Verderio C, Varoqui H, Erickson JD, Matteoli M (2002) Localization and functional relevance of system a neutral amino acid transporters in cultured hippocampal neurons. *J Biol Chem* **277**, 10467–73.
 15. Dolinska M, Zablocka B, Sonnewald U, Albrecht J (2004) Glutamine uptake and expression of mRNA's of glutamine transporting proteins in mouse cerebellar and cerebral cortical astrocytes and neurons. *Neurochem Int* **44**, 75–81.
 16. Ikonomidou C, Qin Y, Labruyere J, Olney JW (1996) Motor neuron degeneration induced by excitotoxin agonists has features in common with those seen in the SOD-1 transgenic mouse model of amyotrophic lateral sclerosis. *J Neuropathol Exp Neurol* **55**, 211–24.
 17. Rothstein JD, Jin L, Dykes-Hoberg M, Kuncl RW (1993) Chronic inhibition of glutamate uptake produces a model of slow neurotoxicity. *Proc Natl Acad Sci USA* **90**, 6591–5.
 18. Casley CS, Land JM, Sharpe MA, Clark JB, Duchon MR, Canevari L (2002) β -Amyloid fragment 25–35 causes mitochondrial dysfunction in primary cortical neurons. *Neurobiol Dis* **10**, 258–67.
 19. Xia Z, Tauskela J, Small DL (2003) Disulfonic stilbenes prevent β -amyloid (25–35) neuronal toxicity in rat cortical cultures. *Neurosci Lett* **340**, 53–6.
 20. Buntup D, Skare O, Solbu TT, Chaudhry FA, Storm-Mathisen J, Thangnipon W (2008) β -amyloid 25–35 peptide reduces the expression of glutamine transporter SAT1 in cultured cortical neurons. *Neurochem Res* **33**, 248–56.
 21. Pike CJ, Burdick D, Walencewicz AJ, Glabe CG, Cotman CW (1993) Neurodegeneration induced by β -amyloid peptides in vitro: the role of peptide assembly state. *J Neurosci* **13**, 1676–87.
 22. Soi-ampornkul R, Tong L, Thangnipon W, Balazs R, Cotman CW (2008) Interleukin-1 β interferes with signal transduction induced by neurotrophin-3 in cultured cortical neurons. *Brain Res* **1188**, 189–97.
 23. Kivell BM, McDonald FJ, Miller JH (2000) Serum-free culture of rat post-natal and fetal brainstem neurons. *Dev Brain Res* **120**, 199–210.
 24. Heckel T, Broer A, Wiesinger H, Lang F, Broer S (2003) Asymmetry of glutamine transporters in cultured neural cells. *Neurochem Int* **43**, 289–98.
 25. Jutapakdeegul N, Casalotti SO, Govitrapong P, Kotchabhakdi N (2003) Postnatal touch stimulation acutely alters corticosterone levels and glucocorticoid receptor gene expression in the neonatal rat. *Dev Neurosci* **25**, 26–33.
 26. Lee BY, Ban JY, Seong YH (2005) Chronic stimulation of GABA_A receptor with muscimol reduces amyloid β protein (25–35)-induced neurotoxicity in cultured rat cortical cells. *Neurosci Res* **52**, 347–56.
 27. Giulietti A, Overbergh L, Valckx D, Decallonne B, Bouillon R, Mathieu C (2001) An overview of real-time quantitative PCR: applications to quantify cytokine gene expression. *Methods* **25**, 386–401.
 28. Baron U, Schnappinger D, Helbl V, Gossen M, Hillen W, Bujard H (1999) Generation of conditional mutants in higher eukaryotes by switching between the expression of two genes. *Proc Natl Acad Sci USA* **96**, 1013–8.
 29. Bolaños JP, Heales SJ, Peuchen S, Barker JE, Land JM, Clark JB (1996) Nitric oxide-mediated mitochondrial damage: a potential neuroprotective role for glutathione. *Free Radic Biol Med* **21**, 995–1001.
 30. Stewart VC, Land JM, Clark JB, Heales SJ (1998) Pretreatment of astrocytes with interferon- α /beta prevents neuronal mitochondrial respiratory chain damage. *J Neurochem* **70**, 432–4.
 31. Heales SJ, Lam AA, Duncan AJ, Land JM (2004) Neurodegeneration or neuroprotection: the pivotal role of astrocytes. *Neurochem Res* **29**, 513–9.
 32. Tian G, Lai L, Guo H, Lin Y, Butchbach ME, Chang Y, Lin CL (2007) Translational control of glial glutamate transporter EAAT2 expression. *J Biol Chem* **282**, 1727–37.
 33. Mattson MP, Cheng B, Davis D, Bryant K, Lieberburg I, Rydel RE (1992) β -Amyloid peptides destabilize calcium homeostasis and render human cortical neurons vulnerable to excitotoxicity. *J Neurosci* **12**, 376–89.
 34. Wei Z, Song MS, MacTavish D, Jhamandas JH, Kar S (2008) Role of calpain and caspase in β -amyloid-induced cell death in rat primary septal cultured neurons. *Neuropharmacology* **54**, 721–33.
 35. Domenici MR, Paradisi S, Sacchetti B, Gaudi S, Balduzzi M, Bernardo A, Ajmone-Cat MA, Minghetti L, et al (2002) The presence of astrocytes enhances beta amyloid-induced neurotoxicity in hippocampal cell cultures. *J Physiol* **96**, 313–6.
 36. Pfrieger FW, Barres BA (1997) Synaptic efficacy enhanced by glial cells in vitro. *Science* **277**, 1684–7.
 37. Dowson VL (1995) Nitric Oxide: role in neurotoxicity. *Clin Exp Pharmacol Physiol* **22**, 305–8.
 38. Peña LA, Brecher CW, Marshak DR (1995) β -amyloid regulates gene expression of glial trophic substances S-100 β in C6 glioma and primary astrocyte cultures. *Mol Brain Res* **34**, 118–26.
 39. Hu J, Akama KT, Krafft GA, Chromy BA, Van Eldik LJ (1998) Amyloid- β peptide activates cultured astrocytes: morphological alterations, cytokine induction and nitric oxide release. *Brain Res* **785**, 195–206.
 40. Fernández-Tomé P, Brera B, Arévalo MA, de Ceballos ML (2004) β -amyloid_{25–35} inhibits glutamate uptake in cultured neurons and astrocytes: modulation of uptake as a survival mechanism. *Neurobiol Dis* **15**, 580–9.
 41. Geci C, How J, Alturaihi H, Kumar U (2007) β -amyloid increases somatostatin expression in cultured cortical neurons. *J Neurochem* **101**, 664–73.